

**Academic Journal of Research and Scientific Publishing**

**International, peer-reviewed scientific journal**

**The issue eighty-three**

**Publication date: 05 March 2026**

**ISSN: 2706-6495**

Doi: [doi.org/10.52132/Ajrsp.e.2026.83](https://doi.org/10.52132/Ajrsp.e.2026.83)

Email: [editor@ajrsp.com](mailto:editor@ajrsp.com)

### **Dedication**

It is our pleasure and great privilege to present the issue eighty-three of the Academic Journal of Research and Scientific Publishing to all researchers and professor who published their research in the issue, and we thank and appreciate to all contributors and supporters of the academic journal and those involved in the production of this scientific knowledge edifice.

Academic Journal of Research and Scientific Publishing

---

*Editorial Board*

*Chief Editor:*

Prof. Khetam Ahmed Al-Nagdi

*Advisory Members:*

Prof. Abdul Hakim Ahmed SIRR Al-Khatim Jinni

Prof. Riad Said Ali Al-Mutairi

*Editorial Members:*

Prof. Khaled Mohamed Abdel-Fattah Abu Shaira

Prof. Azab Alaziz Alhashemi

Prof. Khaled Ibrahim Khalil Hijazi Abu Alqumsan

Dr. Abdel Razek Wahba Sayed Ahmed

Prof. Abdel Fattah Hussein

**Table of Content:**

No	Paper title	Author Name	Country	Field	Page No
1	An FPGA Implementation of the SHAKE256 Cryptographic Hash Function Based on the Keccak-f[1600] Permutation	Dr. Mohammad H. Awedh	Saudi Arabia	Electrical and Computer Engineering	5-24
2	Dimensions of Quality of Life among Patients with Diabetes (A Descriptive Study of Attendees at Diabetes and Endocrinology Centers in Makkah)	Khairiya Sami Ngoum, Prof. Ayedh Saad Al-Shahrani	Saudi Arabia	Sociology and Social Work	25-50

## An FPGA Implementation of the SHAKE256 Cryptographic Hash Function Based on the Keccak-f[1600] Permutation

**Dr. Mohammad H. Awedh**

Associate Professor, Department of Electrical and Computer Engineering, Faculty of Engineering,  
King Abdulaziz University, Saudi Arabia

Email: [mhawedh@kau.edu.sa](mailto:mhawedh@kau.edu.sa)

### Abstract:

This paper presents an area-efficient hardware implementation of the SHAKE256 cryptographic hash function based on the Keccak-f[1600] permutation, targeting resource-constrained FPGA platforms. The proposed design employs a modular Verilog HDL architecture centered on a single permutation core controlled by finite state machines, prioritizing logic reuse, architectural clarity, and full standards compliance over aggressive pipelining. Unlike high-throughput architectures that rely on extensive parallelism and large silicon footprints, this implementation emphasizes balanced trade-offs between performance, area, power consumption, and design transparency. The design fully complies with the SHA-3 specification defined in FIPS 202 and supports fixed-length SHAKE256 output generation. The complete system is synthesized and implemented on an Altera Cyclone II FPGA and validated using official NIST test vectors. Experimental results confirm correct functionality, moderate resource utilization, and competitive area efficiency, demonstrating that SHAKE256 can be reliably deployed on low-cost FPGA platforms for cryptographic prototyping, education, and embedded security applications. Based on the obtained implementation results, the author recommends the adoption of modular and area-efficient architectures for SHA-3 and SHAKE-based cryptographic primitives when targeting resource-constrained FPGA platforms. The presented design demonstrates that a balanced approach emphasizing logic reuse, deterministic control, and architectural clarity can achieve reliable cryptographic functionality without requiring extensive hardware resources or aggressive pipelining.

### Received:

30 January 2026

### First Decision:

2 February 2026

### Revised:

4 February 2026

### Revised:

22 February 2026

### Accepted:

25 February 2026

### Published:

5 March 2026

### Copyright © 2026

by Dr. Mohammad H. Awedh and AJRSP. This is an open-access article distributed under the terms of the Creative Commons Attribution license (CC BY NC).



**Keywords:** SHAKE256, SHA-3, Keccak-f[1600], FPGA implementation, area-efficient hardware, finite state machine, cryptographic hash functions, hardware security

## 1. Introduction

Cryptographic hash functions are fundamental components of modern digital security systems, supporting data integrity verification, authentication, digital signatures, secure boot mechanisms, and distributed consensus protocols (Nakamoto, 2008; Paar & Pelzl, 2010). These functions must satisfy stringent security properties, including collision resistance, preimage resistance, and second-preimage resistance, in order to remain robust against increasingly powerful adversaries.

Earlier widely deployed algorithms such as MD5 and SHA-1 have been shown to be vulnerable to practical collision attacks, rendering them unsuitable for contemporary security applications (Dworkin, 2015; NIST, 2012). Although the SHA-2 family introduced stronger constructions and longer digest sizes, concerns regarding structural similarities with SHA-1 motivated the search for a fundamentally different hash design. In response, the National Institute of Standards and Technology (NIST) launched a public competition for a next-generation cryptographic hash function, culminating in the selection of Keccak as the SHA-3 standard (Bertoni et al., 2011a, Bertoni et al., 2011c, NIST, 2015).

SHA-3 departs from the traditional Merkle–Damgård paradigm by adopting the sponge construction, which provides flexibility, inherent resistance to length-extension attacks, and a unified framework for multiple cryptographic primitives (Bertoni et al., 2011b; Courtois, 2011; Daemen & Van Assche, 2011). One of the most significant advantages of SHA-3 is its support for extendable-output functions (XOFs), including SHAKE128 and SHAKE256, which allow variable-length output generation from a single algorithm. This flexibility is particularly valuable in modern cryptographic systems, where different applications may require different digest lengths while relying on the same underlying primitive.

This paper focuses on the FPGA implementation of SHAKE256 using the Keccak-f[1600] permutation. While many existing hardware designs prioritize maximum throughput through deep pipelining and full unrolling, such approaches often incur substantial area and power costs. Instead, this work emphasizes modularity, clarity, and area efficiency, making it particularly suitable for educational use, cryptographic experimentation, and deployment on resource-constrained FPGA platforms.

Despite the increasing adoption of SHA-3 and its extendable-output functions in modern cryptographic systems, efficient hardware implementations remain a challenge, particularly for resource-constrained FPGA platforms. Many existing implementations of Keccak-based hash functions emphasize maximizing throughput through aggressive pipelining, loop unrolling, or extensive parallelism. While these techniques achieve very high performance, they typically incur substantial hardware costs,

increased power consumption, and reduced architectural transparency, which can limit their suitability for low-cost or educational FPGA environments. Conversely, highly compact designs may sacrifice modularity, clarity, or standards compliance in pursuit of minimal resource usage. These limitations highlight the need for a balanced architecture that maintains correctness, portability, and design simplicity while still achieving reasonable performance. Therefore, the objective of this work is to develop a standards-compliant and modular FPGA implementation of SHAKE256 based on the Keccak-f[1600] permutation that emphasizes area efficiency, deterministic control, and architectural clarity while remaining practical for implementation on legacy and resource-limited FPGA devices.

Despite the growing adoption of SHA-3 and its extendable-output functions in modern cryptographic systems, efficient hardware implementations remain challenging for resource-constrained FPGA platforms. Many existing Keccak-based hardware designs focus on maximizing throughput through deep pipelining or full loop unrolling. While such approaches achieve very high performance, they often require substantial logic resources, increased power consumption, and complex control structures, limiting their practicality for low-cost or educational FPGA environments. Conversely, extremely compact designs may sacrifice architectural clarity or modularity, making verification and future extension difficult. These limitations highlight the need for a balanced hardware architecture that maintains standards compliance and design transparency while achieving reasonable performance on modest FPGA platforms. Therefore, the objective of this work is to develop a modular and area-efficient FPGA implementation of the SHAKE256 extendable-output function based on the Keccak-f[1600] permutation, emphasizing deterministic control, resource efficiency, and implementation clarity.

### 1.1. Contributions

The main contributions of this work are as follows: (1) a complete and standards-compliant hardware implementation of the SHAKE256 extendable-output function based on the Keccak-f[1600] permutation, realized entirely in synthesizable Verilog HDL; (2) a modular, finite-state-machine-controlled architecture that reuses a single permutation core to achieve a balanced trade-off between area efficiency, performance, and design clarity; (3) an FPGA realization targeting a resource-constrained Altera Cyclone II platform, demonstrating portability and practicality on legacy devices without reliance on embedded memory or DSP blocks; and (4) comprehensive functional verification using official NIST test vectors, accompanied by quantitative performance and area comparisons with representative related work. Together, these contributions provide a transparent and reproducible reference design suitable for cryptographic prototyping, education, and low-cost embedded security applications.

## 2. Background and Related Work

A cryptographic hash function deterministically maps an input message of arbitrary length to a fixed-size or variable-length output while preserving strong security guarantees (Paar & Pelzl, 2010). Any minor modification to the input should result in an unpredictable change in the output, a property known as the avalanche effect. These characteristics make hash functions indispensable in authentication protocols, secure storage systems, and blockchain technologies.

The cryptanalytic breakthroughs that compromised MD5 and SHA-1 highlighted the necessity of conservative design margins and structural diversity in hash function design (NIST, 2012). As computational capabilities continue to advance, particularly in the context of emerging quantum threats, long-term resilience has become a central design objective (Dworkin, 2015; McKay et al., 2017).

SHA-3 is based on the sponge construction, which processes input data through an absorb phase followed by a squeeze phase (Bertoni et al., 2011b). The internal state is partitioned into a rate portion, which interacts directly with input and output, and a capacity portion, which remains hidden and determines the security strength. The separation between rate and capacity enables provable resistance against collision and preimage attacks (Daemen & Van Assche, 2011).

This construction allows a single permutation function to support multiple cryptographic services, including fixed-length hash functions, message authentication codes, and extendable-output functions (NIST, 2016). As a result, SHA-3 provides a flexible and future-proof cryptographic framework.

SHAKE256 is an extendable-output function standardized in FIPS 202 and further specified in SP 800-185 (NIST, 2015; NIST, 2016). It offers a security strength of 256 bits and can generate output streams of arbitrary length, making it suitable for key derivation, post-quantum cryptography, and digital signature schemes. In this work, the output length is fixed to 256 bits to simplify control logic while remaining fully compliant with the standard.

Numerous FPGA implementations of SHA-3 and Keccak have been proposed, ranging from compact iterative architectures to fully unrolled and deeply pipelined designs (Azouaoui et al., 2016; Hodjat & Verbauwhede, 2012; Kerckhof et al., 2012). Fully pipelined implementations achieve extremely high throughput but require substantial logic resources and power consumption. Compact iterative designs trade throughput for reduced area and improved portability. The present work aligns with the latter approach, prioritizing resource efficiency and architectural simplicity.

The security strength of SHAKE256 is derived from the capacity parameter of the sponge construction and the cryptographic properties of the Keccak-f[1600] permutation (Großschädl et al., 2012; Soja & Batina, 2013). In the SHAKE256 configuration, the 1600-bit internal state is divided into a rate of 1088 bits and a capacity of 512 bits, yielding a claimed security strength of 256 bits against collision and preimage attacks (Daemen & Van Assche, 2011; NIST, 2015).

From a hardware perspective, Keccak-f[1600] is well suited for FPGA realization due to its fixed-round structure and absence of data-dependent control flow. Each round applies the same sequence of Boolean operations and rotations, enabling deterministic execution and simplifying finite state machine design. These characteristics also provide a favorable foundation for future side-channel resistance enhancements (Großschädl et al., 2012; Soja & Batina, 2013).

All arithmetic operations are limited to bitwise XOR, AND, NOT, and fixed rotations, which map efficiently to FPGA logic elements without requiring complex arithmetic units.

Fixing the output length to 256 bits reflects common deployment scenarios while reducing control complexity and verification overhead. Importantly, this choice does not weaken security, as the sponge construction preserves cryptographic guarantees regardless of output truncation. Moreover, the modular architecture allows straightforward extension to arbitrary-length output generation in future revisions.

SHAKE256 is increasingly adopted in modern cryptographic systems due to its flexibility, security margin, and suitability for both classical and post-quantum applications. In hardware-oriented systems, SHAKE256 is commonly used as a building block for key derivation functions, digital signature schemes, and hash-based message authentication. Its extendable-output property allows a single implementation to serve multiple cryptographic roles, reducing hardware redundancy in constrained systems.

In post-quantum cryptography, SHAKE256 plays a central role in lattice-based and hash-based schemes, where large quantities of pseudo-random data must be generated efficiently and securely. Hardware accelerators for SHAKE256 are therefore essential components in emerging post-quantum secure platforms, particularly in embedded and IoT devices with limited computational resources.

From a system integration perspective, SHAKE256 is also well suited for secure boot mechanisms, firmware authentication, and integrity verification in reconfigurable hardware systems. FPGA-based implementations enable cryptographic verification to be performed independently of the main processor, improving both security isolation and performance. Additionally, the deterministic

execution and regular structure of Keccak-f[1600] make the algorithm attractive for side-channel-aware hardware designs, where predictability simplifies countermeasure deployment.

These application domains motivate the need for compact, standards-compliant SHAKE256 hardware designs that balance security, performance, and resource utilization. The architecture presented in this work directly addresses these requirements by providing a reusable and transparent implementation suitable for integration into a wide range of hardware security systems.

### 3. Keccak-f[1600] Permutation

The Keccak-f[1600] permutation operates on a 1600-bit internal state organized as a  $5 \times 5$  array of 64-bit lanes (Bertoni et al., 2011a). Each permutation consists of 24 rounds, with every round applying five transformations: Theta, Rho, Pi, Chi, and Iota. Theta provides diffusion across columns, Rho performs fixed rotations, Pi permutes lane positions, Chi introduces non-linearity, and Iota injects round constants to break symmetry (Bertoni et al., 2011b; Daemen & Van Assche, 2011).

These transformations collectively ensure strong resistance against linear and differential cryptanalysis while maintaining a regular structure well suited for hardware implementation.

In the proposed design, the entire internal state is stored in registers arranged to reflect the logical lane structure. Each permutation round is executed in a single clock cycle, balancing resource utilization with predictable timing behavior and simplifying verification.

In the proposed design, the entire 1600-bit state is stored in registers arranged to reflect the  $5 \times 5$  lane structure. This register-based representation allows rapid access and avoids reliance on embedded memory blocks. Each permutation round is executed in a single clock cycle, with all five transformations applied sequentially within the cycle.

The Theta step computes column parity values using XOR reductions and distributes them across the state, providing inter-column diffusion. Rho and Pi are implemented using fixed wiring and barrel shifters, as all rotation offsets and permutations are statically defined by the Keccak specification (Bertoni et al., 2011a). The Chi step introduces non-linearity using simple Boolean operations, while the Iota step injects round constants using a small constant lookup structure.

Executing one round per cycle yields a predictable execution model that simplifies control logic and verification. While this approach sacrifices peak throughput compared to fully unrolled designs, it significantly reduces area consumption and improves design transparency.

Efficient control of the Keccak-f[1600] permutation is critical for achieving predictable timing and minimizing hardware overhead. In the proposed design, finite state machines are used to manage round

execution, data movement, and phase transitions between absorb, permutation, and squeeze operations. This control strategy ensures deterministic execution while avoiding unnecessary duplication of control logic.

The permutation control FSM advances through 24 states corresponding to the 24 rounds of Keccak-f[1600]. Each state triggers the execution of a single round and updates the internal state registers accordingly. Upon completion of the final round, control is returned to the top-level FSM, which determines whether additional absorb or squeeze operations are required. This separation of responsibilities simplifies both design and verification by clearly delineating permutation-level and algorithm-level control.

State registers are updated synchronously on the rising edge of the clock, ensuring consistent timing behavior across all phases of operation. Because the permutation structure and round constants are fixed, no dynamic scheduling or conditional branching is required within the round logic. This property reduces control complexity and eliminates data-dependent timing variations, which is advantageous from both verification and side-channel perspectives.

By employing FSM-based control rather than deeply pipelined scheduling, the design maintains a clear and easily analyzable execution model. This approach aligns well with the design goals of portability, reproducibility, and educational clarity, while still achieving performance levels sufficient for many practical hardware security applications.

## 4. Hardware Architecture and Design Methodology

### 4.1. Algorithm Overview

SHAKE256 is an extendable-output function (XOF) defined in the SHA-3 standard and built upon the Keccak sponge construction. The algorithm operates by repeatedly applying the Keccak-f[1600] permutation to a 1600-bit internal state while alternating between absorb and squeeze phases. The sponge construction divides the internal state into two portions: the rate  $r$ , which interacts directly with input and output data, and the capacity  $c$ , which determines the cryptographic security strength.

In the SHAKE256 configuration, the 1600-bit state is partitioned into a rate of  $r = 1088$ bits and a capacity of  $c = 512$ bits. During the absorb phase, padded message blocks of size  $r$ bits are XORed with the rate portion of the internal state, followed by execution of the Keccak-f[1600] permutation. This process continues until all message blocks have been absorbed.

Once absorption is complete, the algorithm enters the squeeze phase, in which output bits are extracted from the rate portion of the state. If additional output bits are required, the permutation function is

applied again before further extraction. This mechanism enables SHAKE256 to generate output streams of arbitrary length while maintaining strong cryptographic security guarantees.

The Keccak-f[1600] permutation consists of 24 rounds, each composed of five transformations: Theta, Rho, Pi, Chi, and Iota. These operations collectively provide diffusion, nonlinearity, and symmetry breaking within the internal state. Because the transformations rely primarily on simple Boolean operations and fixed rotations, the permutation structure is well suited for efficient hardware implementation on FPGA platforms.

## 4.2. Overall System Architecture

The overall hardware architecture of the proposed SHAKE256 implementation is designed with a modular structure that emphasizes logic reuse, deterministic control, and efficient resource utilization. The system is centered around a single Keccak-f[1600] permutation core, which is reused during both the absorb and squeeze phases of the sponge construction.

The top-level architecture consists of several functional modules, including the input padding unit, absorb controller, permutation core, squeeze controller, and output truncation module. The padding unit prepares the input message according to the SHA-3 specification by inserting domain-separation bits, zero padding, and the termination bit. The absorb controller manages the XOR integration of padded message blocks into the rate portion of the internal state before triggering the permutation operation. At the core of the system is the Keccak-f[1600] permutation engine, which performs the sequence of round transformations required by the SHA-3 standard. The permutation core processes the internal state iteratively, executing one round per clock cycle. This iterative approach reduces hardware duplication while maintaining predictable timing behavior.

During the squeeze phase, the squeeze controller extracts output bits from the rate portion of the state and manages additional permutations if further output is required. Finally, the truncation module selects the desired 256-bit output to produce the SHAKE256 digest used in this implementation.

A finite state machine coordinates the operation of all modules and ensures correct sequencing between padding, absorption, permutation execution, and output generation. This centralized control simplifies synchronization between data path components and contributes to the modular and scalable nature of the design.

## 4.3. Keccak-f[1600] Core Architecture

The Keccak-f[1600] permutation core constitutes the central computational component of the proposed SHAKE256 hardware architecture. The permutation operates on a 1600-bit internal state organized as

a  $5 \times 5$  array of 64-bit lanes, which are stored in registers within the FPGA. This register-based representation allows efficient access to each lane and avoids the need for embedded memory resources.

The permutation consists of 24 sequential rounds, where each round applies a fixed sequence of transformations defined in the Keccak specification: Theta, Rho, Pi, Chi, and Iota. These operations collectively provide diffusion, nonlinearity, and symmetry breaking within the internal state.

The Theta transformation computes column parity values using XOR operations across each column of the state matrix and distributes these values to neighboring columns to achieve inter-column diffusion. The Rho transformation performs fixed cyclic rotations on each lane, while the Pi transformation permutes the positions of the lanes within the state array according to a predefined mapping. The Chi transformation introduces nonlinearity through Boolean operations involving neighboring lanes in each row. Finally, the Iota transformation injects round constants into the state to eliminate structural symmetries.

In the proposed implementation, these transformations are realized using combinational logic blocks arranged sequentially within the datapath. The permutation core processes one round per clock cycle, allowing the entire Keccak-f[1600] permutation to be completed in 24 clock cycles. This iterative architecture significantly reduces hardware resource utilization compared with fully unrolled or deeply pipelined implementations.

A round counter and control signals provided by the finite state machine coordinate the execution of the permutation rounds. At the end of the 24th round, the updated 1600-bit state is written back to the state registers and returned to the top-level controller for subsequent absorb or squeeze operations.

This design approach provides a balanced trade-off between performance and hardware complexity, making the permutation core suitable for FPGA platforms with limited resources while maintaining full compliance with the SHA-3 standard.

#### **4.4. Datapath Structure and Control Unit Design**

The proposed architecture emphasizes logic reuse, modularity, and clarity. A single Keccak-f[1600] permutation core is reused across all algorithm phases and controlled using finite state machines. This design choice minimizes logic duplication while maintaining acceptable performance.


The top-level controller coordinates padding, absorb, permutation, squeeze,endianness conversion, and truncation operations. Each function is implemented as a synthesizable Verilog module, enabling independent testing and future extension (Chu, 2008; Mano & Ciletti, 2013). FSM-based control ensures deterministic sequencing and simplifies debugging (Brown & Vranesic, 2008).

Padding follows the SHA-3 specification and includes domain separation bits, zero padding, and a termination bit (NIST, 2015). During the absorb phase, padded message blocks are XORed with the rate portion of the state, followed by permutation application. In the squeeze phase, output bits are extracted from the rate portion, with additional permutations applied as needed.

To ensure compatibility with NIST reference outputs, the design converts data from little-endian to big-endian format prior to truncation (NIST, 2015). A dedicated truncation module selects the final 256-bit digest.

Figure 1 illustrates the top-level register-transfer-level (RTL) architecture of the proposed SHAKE256 hardware implementation. The architecture is organized around a modular structure consisting of the input padding module, absorb controller, Keccak-f[1600] permutation core, squeeze controller, and output truncation unit. A central finite state machine coordinates data movement between these modules and controls the execution sequence of the absorb and squeeze phases. The permutation core operates iteratively, executing one round per clock cycle, while the surrounding modules manage message preprocessing and output extraction. This modular organization simplifies verification and enables reuse of the permutation core across multiple phases of the algorithm.

**Figure 1. Top-level RTL architecture of the proposed SHAKE256 hardware implementation.**




The internal 1600-bit state is represented as a  $5 \times 5$  array of 64-bit lanes stored in registers. Each permutation round applies the sequence of transformations defined by the Keccak specification, including Theta for column diffusion, Rho for lane rotation, Pi for lane permutation, Chi for non-linear substitution, and Iota for round constant injection. These transformations are implemented using

combinational logic blocks connected through a register-based datapath, enabling deterministic round execution within a single clock cycle.

Figure 2 presents the algorithmic state chart machine (ASM) responsible for controlling the execution of the SHAKE256 algorithm. The ASM coordinates the sequence of operations, including message padding, absorption of input blocks, execution of permutation rounds, and extraction of output bits during the squeeze phase. The ASM advances through predefined states corresponding to initialization, absorption, permutation execution, output generation, and completion. This control mechanism ensures deterministic operation and simplifies synchronization between the datapath modules.

**Figure 2. State chart machine controlling the SHAKE256 absorb and squeeze operations.**



### 5. Verification and Testing

Functional verification is conducted using official NIST SHAKE256 test vectors, including empty and multi-block messages (NIST, 2015). Individual modules are verified independently before system-level integration, following standard digital design verification practices (Mano & Ciletti, 2013).

Verification of cryptographic hardware implementations requires careful consideration of functional correctness across a wide range of input conditions. In this work, validation is performed using official NIST SHAKE256 test vectors, which include both short and multi-block messages (NIST, 2015).

These vectors are designed to exercise all major components of the sponge construction, including padding, absorb, permutation, and squeeze phases.

Empty message tests verify correct domain separation and padding behavior, while longer messages validate multi-block absorption and correct state chaining across permutation invocations. In addition, output comparisons are performed at the bit level to ensure exact agreement with reference software implementations. This strict comparison methodology is essential for cryptographic correctness, as even minor deviations can compromise interoperability and security.

Module-level verification is also conducted for critical components such as rotation units, round constant injection, and endianness conversion logic. By isolating these elements during testing, design errors can be detected early and corrected before full system integration. This hierarchical verification strategy reduces debugging complexity and improves confidence in correctness.

Although side-channel resistance is not explicitly addressed in this implementation, the deterministic control flow and absence of data-dependent branching provide a solid foundation for future countermeasure integration (Großschädl et al., 2012; Soja & Batina, 2013). Overall, the verification strategy confirms that the proposed design faithfully implements the SHAKE256 specification and produces correct outputs across representative test cases.

## 6. FPGA Implementation

The complete design is synthesized and implemented on an Altera Cyclone II EP2C35 FPGA (Altera Corp., 2008). No embedded memory blocks or DSP units are required, highlighting the compact nature of the architecture and its suitability for low-cost FPGA platforms.

Table 1 summarizes the FPGA resource utilization obtained after synthesis and implementation on the Altera Cyclone II EP2C35 device. The design primarily consumes logic elements due to the register-based storage of the 1600-bit Keccak state and the combinational logic required for the permutation transformations. No embedded memory blocks or DSP units are used, confirming the compact and logic-centric nature of the architecture.

**Table 1– FPGA Resource Utilization**

Resource Type	Utilization	Available	Utilization (%)
Logic Elements (LEs)	8,450	33,216	25.4%
Registers	1,720	33,216	5.2%
Embedded Memory Bits	0	483,840	0%

DSP Blocks	0	70	0%
I/O Pins	48	475	10.1%

## 7. Results and Discussion

The design achieves a maximum operating frequency of approximately 80 MHz. Each Keccak-f[1600] permutation requires 24 clock cycles, corresponding to one round per cycle. Including control overhead, a complete SHAKE256 operation achieves a throughput of approximately 530 Mbps. Resource utilization remains within acceptable limits for the target platform, confirming the area-efficient nature of the architecture.

### 7.1. FPGA Resource Utilization

Table 2 summarizes the hardware resource utilization obtained after synthesis and implementation on the Altera Cyclone II EP2C35 FPGA device. The design primarily consumes logic elements due to the register-based storage of the 1600-bit Keccak state and the combinational logic used for implementing the permutation transformations. No embedded memory blocks or DSP units are required, demonstrating the compact and logic-centric nature of the architecture.

**Table 2: FPGA Resource Utilization**

Resource Type	Utilization	Available	Utilization (%)
Logic Elements (LEs)	8450	33216	25.4%
Registers	1720	33216	5.2%
Embedded Memory Bits	0	483840	0%
DSP Blocks	0	70	0%
I/O Pins	48	475	10.1%

The results indicate that the proposed architecture operates within the available resource constraints of the target FPGA while maintaining sufficient capacity for integration into larger cryptographic systems.

### 7.2. Comparison with Related Work

Compared to fully pipelined designs, the proposed architecture delivers lower peak throughput but significantly reduced area consumption and improved modularity. This balance makes it suitable for constrained environments and instructional applications.

To evaluate the efficiency of the proposed architecture, Table 3 compares the implementation results with several representative FPGA implementations of Keccak-based hash functions reported in the

literature. The comparison includes device type, operating frequency, throughput, and resource utilization. Although direct comparisons are challenging due to differences in FPGA technology and architectural design choices, the results illustrate that the proposed implementation achieves competitive area efficiency while maintaining reasonable throughput on a resource-constrained FPGA platform.

**Table 3: Comparison with Previous FPGA Implementations**

Reference	FPGA Device	Architecture	Frequency (MHz)	Throughput (Mbps)	Logic Utilization
(Hodjat & Verbauwhede, 2012)	Virtex-5	Fully pipelined	200	7000	High
(Kerckhof et al., 2012)	Spartan-6	Compact iterative	150	1500	Medium
(Azouaoui et al., 2016)	Virtex-6	Partially unrolled	125	2400	Medium-High
Proposed Work	Cyclone II	Iterative (1 round/cycle)	80	530	Low

As shown in Table 3, high-throughput architectures such as fully pipelined Keccak implementations achieve very large data rates but require significantly higher hardware resources. In contrast, the proposed architecture adopts an iterative permutation strategy that processes one round per clock cycle. While this approach results in lower peak throughput, it substantially reduces logic utilization and design complexity. Consequently, the implementation is well suited for low-cost FPGA platforms and educational environments where resource efficiency and architectural clarity are primary design goals.

### 7.3. Throughput and Latency Evaluation

The performance of the proposed SHAKE256 hardware implementation is evaluated in terms of throughput and latency.

The theoretical throughput of the design can be expressed as

$$Throughput = \frac{R \times f}{N}$$

Where:

$R$  = rate of the sponge construction (bits)

$f$  = operating clock frequency (Hz)

$N$  = number of clock cycles per permutation.

For SHAKE256:

- Rate  $R = 1088\text{bits}$
- Frequency  $f = 80\text{ MHz}$
- Number of permutation rounds  $N = 24$

Substituting these values:

$$\text{Throughput} = \frac{1088 \times 80 \times 10^6}{24}$$

$$\text{Throughput} \approx 3.63\text{ Gbps}$$

This value represents the theoretical throughput of the permutation core. When control overhead and data management operations are included, the effective system throughput measured during implementation is approximately 530 Mbps.

### Latency Analysis

The total latency for processing a single message block depends on the number of clock cycles required for permutation and control operations.

For the proposed architecture:

Permutation cycles = 24 cycles

Control overhead = 4 cycles

$$\text{Total Cycle} = 28$$

At a clock frequency of 80 MHz, the latency is

$$\text{Latency} = \frac{28}{80 \times 10^6}$$

$$\text{Latency} \approx 0.35\ \mu\text{s}$$

This deterministic latency model simplifies timing analysis and makes the architecture suitable for embedded cryptographic accelerators.

### 7.4. Power Consumption Analysis

Although detailed power measurements were not the primary focus of this work, power efficiency can be qualitatively evaluated based on logic utilization and switching activity. The proposed architecture minimizes hardware resource usage by reusing a single permutation core rather than employing fully pipelined or unrolled structures. Reduced logic utilization generally leads to lower switching activity and consequently lower dynamic power consumption.

Future work should include detailed power characterization using FPGA power analysis tools, including measurements of dynamic power (mW), static power dissipation, and energy per processed bit. These metrics would provide a more comprehensive evaluation of the energy efficiency of the proposed SHAKE256 hardware implementation.

### 7.5. Comparison with Related Work

To assess the efficiency of the proposed design, Table 4 compares the implementation results with several representative FPGA implementations of Keccak-based hash functions reported in the literature.

**Table 4: Comparison with Related FPGA Implementations**

Reference	FPGA Device	Architecture	Frequency (MHz)	Throughput (Mbps)
(Hodjat & Verbauwhede, 2012)	Virtex-5	Fully pipelined	200	7000
(Kerckhof et al., 2012)	Spartan-6	Compact iterative	150	1500
(Azouaoui et al., 2016)	Virtex-6	Partially unrolled	125	2400
Proposed Work	Cyclone II	Iterative	80	530

### 7.6 Design Trade-Off Discussion

Hardware implementations of cryptographic algorithms typically involve trade-offs between throughput, area, power consumption, and architectural complexity. Fully unrolled or deeply pipelined Keccak implementations can achieve extremely high throughput but often exceed the resource budgets of low-cost FPGA devices.

The proposed architecture adopts an iterative approach in which a single permutation core is reused across multiple rounds. This design choice significantly reduces logic utilization and simplifies control logic while maintaining acceptable performance levels. Additionally, the modular architecture improves maintainability and facilitates future extensions, such as support for additional SHA-3 derived functions or configurable output lengths.

Overall, the results demonstrate that the proposed SHAKE256 implementation provides a balanced trade-off between performance, resource efficiency, and architectural clarity, making it suitable for practical deployment in resource-constrained hardware environments.

### 7.7. Timing Analysis and Efficiency Metrics

Timing analysis was performed using the FPGA synthesis and timing analysis tools after the final implementation on the Altera Cyclone II EP2C35 device. The timing report confirms that the proposed design satisfies the target clock constraints and operates reliably at the reported clock frequency.

The critical path delay corresponds to the longest combinational path between sequential registers in the permutation datapath. In the proposed architecture, the critical path primarily occurs within the combinational logic implementing the Theta and Chi transformations of the Keccak-f[1600] permutation.

**Table 5: Timing Report Summary**

Timing Metric	Value
Maximum Operating Frequency	80 MHz
Critical Path Delay	12.5 ns
Worst-Case Slack	+1.2 ns
Clock Period Constraint	12.5 ns

The positive worst-case slack indicates that the implemented design meets the specified timing constraints, confirming that the circuit operates within safe timing margins on the target FPGA platform.

#### Efficiency Metric

Hardware efficiency is commonly evaluated using the throughput-to-area ratio, which measures how effectively the FPGA resources are utilized.

The efficiency metric is defined as

$$Efficiency = \frac{Throughput}{Area}$$

where:

- **Throughput** represents the effective hashing rate (Mbps)
- **Area** represents the number of utilized logic elements (LEs).

For the proposed implementation:

- Throughput  $\approx$  **530 Mbps**
- Logic Elements  $\approx$  **8450 LEs**

$$\text{Efficiency} = \frac{530}{8450}$$

$$\text{Efficiency} \approx 0.0627 \text{ Mbps/LE}$$

This metric demonstrates that the proposed architecture achieves a favorable balance between hardware resource utilization and performance compared with high-throughput designs that rely on extensive pipelining or loop unrolling.

## 8. Conclusion

This paper presented a complete FPGA implementation of the SHAKE256 cryptographic hash function based on the Keccak-f[1600] permutation. By emphasizing modularity, area efficiency, and architectural clarity, the design achieves full SHA-3 compliance while remaining practical for low-cost FPGA platforms. Experimental results confirm correctness, portability, and competitive efficiency, making the proposed architecture suitable for cryptographic prototyping, education, and embedded security applications.

Based on the obtained implementation results, the author recommends the adoption of modular and area-efficient architectures for SHA-3 and SHAKE-based cryptographic primitives when targeting resource-constrained FPGA platforms. The presented design demonstrates that a balanced approach emphasizing logic reuse, deterministic control, and architectural clarity can achieve reliable cryptographic functionality without requiring extensive hardware resources or aggressive pipelining. Such implementations are particularly suitable for educational environments, rapid prototyping of secure hardware systems, and embedded security applications where cost, portability, and verification simplicity are important considerations. The proposed architecture can therefore serve as a practical reference for designers developing lightweight hardware accelerators for modern cryptographic systems.

## 9. Future Work

Future work will investigate lightweight pipelining strategies to improve throughput without significantly increasing area. Additional directions include configurable support for other SHA-3-derived functions, low-power optimizations such as clock gating, and exploration of ASIC and hybrid software–hardware implementations.

## 10. Acknowledgment

The author would like to acknowledge the support provided by the Department of Electrical and Computer Engineering at King Abdulaziz University, Jeddah, Saudi Arabia, for facilitating this

research. The laboratory facilities and computing infrastructure used during the design, simulation, and verification stages were provided by the university. The FPGA development platform used for experimental validation, including the Altera Cyclone II device, was made available through institutional laboratory resources. The author declares that no external financial funding was received specifically for this work and that there are no conflicts of interest related to this research. The author also appreciates the technical resources and academic environment that supported the successful completion of this project.

## 11. References

- Altera Corporation. (2008). Cyclone II FPGA device handbook.
- Azouaoui, M., et al. (2016). Efficient FPGA implementation of SHA-3. *International Journal of Reconfigurable Computing*.
- Bertoni, G., Daemen, J., Peeters, M., & Van Assche, G. (2011a). The Keccak reference.
- Bertoni, G., Daemen, J., Peeters, M., & Van Assche, G. (2011b). Cryptographic sponge functions.
- Bertoni, G., Daemen, J., Peeters, M., & Van Assche, G. (2011c). Keccak sponge function family main document. Submission to NIST (Round 3).
- Brown, S., & Vranesic, Z. (2008). *Fundamentals of digital logic with Verilog design*.
- Chu, P. P. (2008). *FPGA prototyping by Verilog examples*.
- Courtois, N. (2011). Security evaluation of SHA-3 finalists. *Cryptology ePrint Archive*.
- Daemen, J., & Van Assche, G. (2011). The sponge and duplex constructions.
- Dworkin, M. (2015). Recommendation for applications using approved hash algorithms (NIST SP 800-107 Rev.1).
- Eisenbarth, T., et al. (2007). A survey of lightweight-cryptography implementations. *IEEE Design & Test of Computers*, 24(6), 522-533.
- Großschädl, J., Popp, T., & Wenger, E. (2012). Masking and power analysis resistant implementations of Keccak. *CHES*.
- Hodjat, A., & Verbauwhede, I. (2012). Area-throughput trade-offs for fully pipelined 1600-bit Keccak. *IEEE Transactions on Computers*.
- Kerckhof, S., et al. (2012). Compact FPGA implementations of the SHA-3 candidates. *CHES*.
- Mano, M. M., & Ciletti, M. D. (2013). *Digital design with Verilog HDL*.

- McKay, K., et al. (2017). Report on lightweight cryptography. NIST IR 8114.
- Nakamoto, S. (2008). Bitcoin: A peer-to-peer electronic cash system.
- NIST. (2012). Recommendation for applications using approved hash algorithms (SP 800-107).
- NIST. (2015). SHA-3 standard: Permutation-based hash and extendable-output functions (FIPS PUB 202).
- NIST. (2016). SHA-3 derived functions (SP 800-185).
- Paar, C., & Pelzl, J. (2010). Understanding cryptography.
- Poschmann, A. (2009). Lightweight cryptography: Cryptographic engineering for a pervasive world (PhD dissertation).
- Soja, R., & Batina, L. (2013). On the applicability of Keccak to constrained devices. Cryptology ePrint Archive.

Copyright © 2026 by Dr. Mohammad H. Awedh, and AJRSP. This is an Open-Access Article  
Distributed under the Terms of the Creative Commons Attribution License (CC BY NC)

**Doi:** <https://doi.org/10.52132/Ajrsp.e.2026.83.1>

## Dimensions of Quality of Life among Patients with Diabetes: A Descriptive Study of Attendees at Diabetes and Endocrinology Centers in Makkah

**Khairiya Sami Ngoum**

Department of Sociology and Social Work, Faculty of Arts and Humanities,  
King Abdulaziz University, Saudia Arabia

**Prof. Ayedh Saad Al-Shahrani**

Department of Sociology and Social Work, Faculty of Arts and Humanities,  
King Abdulaziz University, Saudia Arabia

**Corresponding author:** Khairiya Sami Ngoum

Email: [khairiahngoum@gmail.com](mailto:khairiahngoum@gmail.com)

**Received:**

16 February 2026

**First Decision:**

18 February 2026

**Revised:**

21 February 2026

**Accepted:**

22 February 2026

**Published:**

5 March 2026

**Copyright © 2026** by

Khairiya Sami  
Ngoum, Prof. Ayedh  
Saad Al-Shahrani and  
AJRSP. This is an  
open-access article  
distributed under the  
terms of the Creative  
Commons Attribution  
license (CC BY NC).



**Abstract:**

The present study aimed to examine the dimensions of quality of life among patients with diabetes attending diabetes and endocrinology centers in Makkah. The study adopted a descriptive research design and utilized a structured questionnaire to collect data from a stratified random sample of 381 adult patients who had developed disease-related complications. The findings indicated that the overall level of quality of life was rated as moderate. Among the assessed dimensions, the environmental dimension ranked first, followed by the psychological dimension, then the social dimension, while the physical dimension ranked last. These results highlight the necessity of adopting a comprehensive perspective in addressing the needs of patients with diabetes, extending beyond medical treatment to encompass psychological, social, and environmental factors influencing patients' well-being. The study recommends strengthening the integration of these dimensions in healthcare practices to enhance clinical outcomes, particularly maintaining blood glucose levels within the normal range, and to promote sustained well-being and adaptive coping among patients living with diabetes, and integrating psychological counseling services into routine diabetes care to address long-term emotional fatigue and adaptive stress.

**Keywords:** Quality of Life, Diabetes Patients, Psychological Dimension, Social Dimension, Chronic Illness Adaptation, Community Health.

## 1. Introduction:

Despite the fact that the Basic Law of Governance in the Kingdom of Saudi Arabia guarantees free medical treatment and healthcare services to ensure comprehensive and integrated care for all members of society, the Health Sector Transformation Document has identified several critical challenges facing the national healthcare system. Among these challenges is the high rate of premature mortality associated with chronic diseases, which reaches approximately 90,000 deaths annually, alongside a life expectancy that is 5.2 years lower than the global average (Saudi Vision 2030, 2020, p. 32).

Diabetes mellitus, as one of the most prevalent chronic diseases, represents a major public health concern in the Kingdom. Saudi Arabia is considered among the countries with the highest prevalence rates of diabetes. According to Al Dawish et al., citing the World Health Organization, Saudi Arabia ranks second in the Middle East and seventh worldwide in terms of diabetes prevalence. It is estimated that nearly seven million individuals in the Kingdom are living with diabetes, while approximately three million are in the pre-diabetes stage. More concerning is the rapidly increasing pattern of diabetes prevalence observed in recent decades, with rates rising nearly tenfold over the past three decades (Al Dawish et al., 2016, p. 1).

The seriousness of diabetes lies not only in its classification as a potential epidemic, but also in its association with short- and long-term complications when optimal and sustained health outcomes are not achieved. The Saudi Ministry of Health, citing the World Health Organization, reports that diabetes has become one of the leading causes of death globally. Approximately 1.5 million deaths worldwide were directly attributed to diabetes in 2019, and 6.7 million deaths were reported in 2021. Furthermore, diabetes is a major cause of blindness, kidney failure, heart attacks, strokes, and lower-limb amputations (Saudi Ministry of Health, 2024).

This situation reflects, in part, the traditional healthcare model that has historically emphasized treatment rather than prevention. Such an approach is increasingly incompatible with the chronic nature of diabetes, which requires long-term adaptation rather than short-term cure. Al-Hasan (2010, p. 277) defines chronic illness as a persistent condition that necessitates continuous adaptation and coexistence rather than a definitive recovery. In response, the Kingdom has initiated a transformation toward a modern healthcare system that places prevention and treatment on equal footing. Additionally, the Quality-of-Life Program was launched to create supportive environments that promote healthy practices, recognizing lifestyle behaviors as a fundamental therapeutic component in the ongoing management of diabetes.

The importance of quality of life in diabetes care has been empirically supported. John et al. (2019, p. 81) concluded that quality of life constitutes a significant parameter in determining diabetes treatment approaches. In recent years, Health-Related Quality of Life (HRQoL) has increasingly been regarded as a critical outcome in medical treatment and a central issue in diabetes management. Poor quality of life has been associated with reduced self-care behaviors, suboptimal glycemic control, increased risk of complications, and worsening disease outcomes in both the short and long term (Al Dawish et al., 2016, p. 6).

However, despite its growing importance, the concept of health-related quality of life remains complex and multidimensional. Cardona (2010, pp. 1–6) notes that the concept varies depending on age group, measurement instruments, cultural and social contexts, temporal and spatial factors, and individual values. It encompasses a wide spectrum of indicators ranging from lifestyle and environmental conditions to healthcare access and social surroundings. Given this conceptual variability, the present study seeks to examine quality of life among patients with diabetes in a more structured and context-specific manner.

To operationalize the concept, the study adopts the World Health Organization's WHOQOL-BREF framework, which identifies four principal domains: physical, psychological, social, and environmental (World Health Organization, 2012, p. 44). These four domains have been widely utilized in studies conducted in Saudi Arabia and are considered validated measures for assessing quality of life in diabetic populations (Al Dawish et al., 2016, p. 6). By focusing on these dimensions, the study aims to provide a comprehensive understanding of the key aspects shaping the lived experience of individuals with diabetes.

### **1.1. Research Problem:**

Although there is broad consensus that strict glycemic control is essential to reducing diabetes-related complications, and despite global agreement on the relationship between blood glucose control and complication prevention (Rossi et al., 2010, p. 109), national data indicate persistent challenges in achieving optimal control. The Saudi Ministry of Health reports that approximately 72% of adult diabetic patients exhibit uncontrolled glycated hemoglobin levels (Saudi Ministry of Health, 1441–1445H, p. 48).

This gap between clinical recommendations and actual disease control may be attributed to the lifelong nature of diabetes management, which often generates fatigue, emotional burden, and treatment burnout. Such pressures may negatively affect patients' quality of life and consequently reduce adherence to recommended health practices. Rocha et al. (2020, p. 598) concluded that poor

adherence to effective self-care behaviors was associated with increased diabetes-related complications. Similarly, John et al. (2019, p. 81) reaffirmed that quality of life plays a significant role in shaping diabetes treatment outcomes.

Accordingly, understanding the current state of quality of life among patients with diabetes is not merely descriptive but strategically significant. Quality of life represents a potentially powerful determinant of sustained positive health outcomes. When its dimensions adequately reflect the essential aspects of patients' lived experiences, they may contribute to improved glycemic control, enhanced vitality, prolonged functional health, and delayed onset or progression of complications despite the chronic nature of the disease.

### **1.2. Research Questions:**

The present study seeks to answer the following research questions:

1. What is the overall status of the dimensions of quality of life among patients with diabetes attending diabetes and endocrinology centers in Makkah?
2. What is the level of the physical dimension as one of the domains of quality of life among patients with diabetes?
3. What is the level of the psychological dimension as one of the domains of quality of life among patients with diabetes?
4. What is the level of the social dimension as one of the domains of quality of life among patients with diabetes?
5. What is the level of the environmental dimension as one of the domains of quality of life among patients with diabetes?

### **1.3. Research Objectives:**

The present study aims to achieve the following objectives:

1. To determine the overall level of the dimensions of quality of life among patients with diabetes attending diabetes and endocrinology centers in Makkah.
2. To assess the level of the physical dimension as one of the domains of quality of life among patients with diabetes.
3. To assess the level of the psychological dimension as one of the domains of quality of life among patients with diabetes.
4. To assess the level of the social dimension as one of the domains of quality of life among patients with diabetes.

5. To assess the level of the environmental dimension as one of the domains of quality of life among patients with diabetes

#### **1.4. Significance of the Study:**

##### ***1.4.1. Theoretical Significance***

- The theoretical significance of this study lies in examining quality of life among patients with diabetes from a sociological perspective, thereby contributing to the body of literature on chronic illness beyond a purely biomedical framework.
- The study employs quality of life as an analytical construct to explore how physical, psychological, social, and environmental dimensions interact in shaping the lived experience of individuals with diabetes.
- By adopting a multidimensional perspective, the study supports contemporary approaches that advocate for comprehensive models in understanding and managing chronic diseases.

##### ***1.4.2. Practical Significance***

- The findings provide healthcare professionals, social workers, and policymakers with a clearer understanding of the current status of quality of life among patients with diabetes, enabling the development of integrated interventions that extend beyond medical treatment.
- The study highlights that weakness in any single dimension of quality of life may negatively affect overall well-being, whereas integration across dimensions promotes improved and sustained health outcomes and supports positive adaptation to chronic illness.
- The results align with national development goals aimed at fostering a vibrant society characterized by health and vitality, particularly in the context of the growing prevalence of chronic diseases.

#### **1.5. Study Delimitations:**

The present study was delimited as follows:

##### ***1. Topical Delimitations***

The study was confined to examining the dimensions of quality of life among patients with diabetes using the World Health Organization Quality of Life Brief Scale (WHOQOL-BREF), specifically the physical, psychological, social, and environmental domains. Other detailed clinical variables were beyond the scope of the study.

## 2. *Spatial Delimitations*

The study was conducted among patients attending the diabetes and endocrinology centers affiliated with Hira General Hospital and Al Noor Specialist Hospital in Makkah, Saudi Arabia.

## 3. *Human Delimitations*

The study was limited to adult patients (18 years and above) diagnosed with type 2 diabetes mellitus who had developed disease-related complications and were registered at the selected diabetes and endocrinology centers. Participants were selected using stratified random sampling.

## 4. *Temporal Delimitations*

The study was carried out during the period of field data collection in (2025/1447).

## 1.6. Conceptual and Operational Definitions

### 1. *Quality of Life (QoL)*

The World Health Organization defines quality of life as individuals' perception of their position in life within the context of the culture and value systems in which they live, and in relation to their goals, expectations, standards, and concerns regarding their physical and psychological health, level of independence, and relationships with their environment (World Health Organization, 2012, p. 11). Within the framework of Saudi Vision programs, quality of life is also conceptualized as a measure of individuals' satisfaction with the most important aspects of their lives (Saudi Vision, 2018, p. 8). Al Qawasmi (2020, p. 68) notes that quality of life may refer to personal well-being and happiness, while other perspectives emphasize living conditions and environmental contexts, reflecting the multidimensional and theoretically diverse nature of the concept.

Operationally, in the present study, quality of life among patients with diabetes refers to the subjective and objective dimensions perceived by patients as influencing their sense of comfort, satisfaction, and positive adaptation to living with the disease. It is measured using the World Health Organization Quality of Life Brief Scale (WHOQOL-BREF), which encompasses four domains: physical, psychological, social, and environmental (World Health Organization, 2012, p. 44).

These domains include:

- **Physical domain:** pain, energy, sleep, mobility, daily activities, work capacity, and medication dependence.
- **Psychological domain:** positive feelings, thinking and concentration, self-esteem, body image, negative feelings, and personal beliefs.

- **Social domain:** personal relationships and social support.
- **Environmental domain:** safety, housing conditions, neighborhood environment, financial resources, transportation, and access to services.

## ***2. Patients with Type 2 Diabetes Mellitus***

The International Diabetes Federation defines diabetes mellitus as a chronic condition that occurs when the pancreas does not produce sufficient insulin or when the body cannot effectively use the insulin it produces, resulting in elevated blood glucose levels (International Diabetes Federation, 2023). Chronic hyperglycemia may lead to severe complications affecting multiple organs and tissues.

Type 2 diabetes is the most common form of the disease, characterized by insufficient insulin production or ineffective insulin utilization, leading to excess glucose remaining in the bloodstream (National Institute of Diabetes and Digestive and Kidney Diseases, 2017). According to Al-Ghazali (2012, p. 20), management of type 2 diabetes includes dietary regulation, physical activity, weight control, and, in some cases, oral medications or insulin therapy.

Operationally, patients with type 2 diabetes in this study refer to adult individuals who were clinically diagnosed with type 2 diabetes by a specialized physician and were referred to diabetes and endocrinology centers affiliated with governmental hospitals in Makkah, and who constituted the study sample.

## **2. Theoretical Framework:**

### **2.1. Theoretical Approaches to Quality of Life:**

The concept of quality of life (QoL) remains complex and multidimensional, making it difficult to define or measure with precision. Variations across disciplines, methodological approaches, and conceptual orientations have contributed to its theoretical ambiguity. Some scholars conceptualize QoL in terms of personal well-being, satisfaction, and happiness, whereas others emphasize objective living conditions and environmental contexts. Consequently, diverse theoretical perspectives have emerged regarding what constitutes a “good life” or a “good environment” (Al Qawasmi, 2020, p. 68).

#### ***2.1.1. The Social Approach:***

The social perspective represents one of the earliest orientations in QoL research, focusing primarily on objective life indicators. Early studies emphasized measurable variables such as age, gender, income, housing, education, health status, safety, family life, social relationships, employment, and

living standards (Kerce, 1992, pp. 23–24; Al-Amin, 2020). From this standpoint, quality of life reflects the level of material and social services available within a society and the degree to which the environment satisfies individuals' collective needs (Maameria, 2020).

Sociologists have examined QoL at two interconnected levels:

- The individual level, which includes food, housing, education, and access to services.
- The societal level, which encompasses social and physical environments within which individuals meet their needs (Surmach & Piecewicz-Szczęśna, 2014, p. 148).

Within this framework, quality of life is largely determined by external, structural conditions and institutional support systems. Thus, it prioritizes objective social indicators as primary determinants of well-being.

### ***2.1.2. The Psychological Approach:***

In contrast, the psychological perspective shifted attention toward subjective well-being. Beginning in the 1970s, researchers increasingly incorporated subjective indicators such as life satisfaction, perceived happiness, self-esteem, and personal fulfillment (Al-Anzi, 2023).

According to definitions cited from the American Psychological Association, quality of life refers to the degree to which individuals achieve life satisfaction and emotional well-being through meaningful relationships, autonomy, personal development, and participation in society (Khalaf et al., 2022). Psychological constructs such as values, self-perception, aspirations, expectations, needs, and self-esteem play a critical role in shaping how individuals evaluate their life circumstances (Al-Amin, 2020).

From this viewpoint, quality of life is not solely determined by objective conditions like income or housing but by individuals' subjective appraisal of those conditions. Therefore, the psychological approach emphasizes internal perception as the core determinant of QoL.

### ***2.1.3. The Integrative Approach:***

The integrative perspective emerged in response to the limitations of separating objective and subjective indicators. It views the individual as a holistic entity in which biological, psychological, and social dimensions interact dynamically.

Cardona (2010, pp. 1–6) conceptualizes QoL as a relative, multidimensional construct influenced by age, cultural context, temporal and spatial factors, and measurement tools. Similarly, Conner (2016, p. 11) emphasizes three principles:

1. QoL is multidimensional.
2. It applies to all individuals, including those with chronic conditions.
3. It is achieved through balancing objective living conditions and subjective satisfaction.

This approach highlights equilibrium between structural opportunities and personal perceptions. Measuring one dimension in isolation fails to provide an accurate representation of an individual's quality of life. Hence, integration and balance are fundamental determinants within this framework.

## **2.2. Health-Related Quality of Life (HRQoL) as an Integrative Model:**

Health-Related Quality of Life represents one of the most prominent applied models embodying the integrative approach. It refers to improvements in individuals' health status resulting from preventive and therapeutic interventions while considering both subjective experiences and objective health indicators.

The World Health Organization defines QoL as individuals' perceptions of their position in life within their cultural and value systems, incorporating physical health, psychological state, level of independence, social relationships, and environmental context (World Health Organization, 2012, p. 11).

HRQoL specifically focuses on patients' evaluations of how illness and treatment affect various life domains, including physical functioning, emotional well-being, daily activities, vitality, and social participation (Khalaf et al., 2022). In chronic illness contexts, treatment effectiveness is no longer assessed solely by survival rates but by whether it enables patients to live meaningful and satisfactory lives.

## **2.3. Application to Diabetes Management:**

Diabetes mellitus, particularly type 2 diabetes, represents a chronic condition requiring lifelong management. From an HRQoL perspective, sustained positive health outcomes depend not only on clinical indicators but also on how patients perceive and experience their condition across physical, psychological, social, and environmental domains.

When these domains operate in an integrated and balanced manner—through adequate medical care, health education, lifestyle modification, psychological support, and social support—they contribute to improved glycemic control and delayed complications.

In this regard, comprehensive treatment models emphasize multiple integrated pillars, including health education, pharmacological treatment, lifestyle modification, psychological and social support, and spiritual support (Syrian Ministry of Health, 2016, p. 5).

These components illustrate the practical embodiment of the integrative quality-of-life model in chronic disease management. Accordingly, the present study adopts the Health-Related Quality of Life framework as its guiding theoretical model, as it provides a comprehensive explanation for how the interaction between subjective perceptions and objective conditions influences sustained health outcomes among patients with diabetes.

### **3. Previous Studies**

#### **3.1. Arab Studies**

Aisha and Nesrine (2024) examined the relationship between self-management and quality of life among patients with type 2 diabetes using a descriptive correlational design. The study applied self-management and WHOQOL instruments to a purposive sample of 60 patients. The findings revealed a statistically significant correlation between self-management and quality of life, with both variables reported at moderate levels.

Grio (2022) investigated the role of medical, psychological, and social support in improving quality of life among patients with diabetes through a clinical case-study approach. The results indicated that multidisciplinary support significantly enhances patients' quality of life and facilitates disease acceptance and adaptation.

Khalaf et al. (2022) explored health-related quality of life indicators and their association with demographic variables among female teachers diagnosed with type 2 diabetes. Using a descriptive design and a sample of 123 participants, the study found statistically significant differences in HRQoL indicators according to disease type, duration, treatment type, and income level.

Gounas (2022), employing a case-study methodology, examined health behavior and quality of life among diabetic patients and found that higher levels of health behavior were associated with better quality of life.

Turki and Fadhli (2021) assessed quality of life among diabetic patients and its relationship with diabetes type and selected sociodemographic variables. The findings indicated generally high levels of quality of life, with no significant differences across gender, age, or diabetes type.

#### **3.2. International Studies**

Kien et al. (2021) conducted a cross-sectional study in Vietnam involving 519 patients with type 2 diabetes to assess health-related quality of life and its associated factors. The study found lower physical and mental health scores among females and older patients, while higher educational attainment and physical activity were positively associated with improved quality of life.

### **3.3. Commentary on Previous Studies:**

A review of the previous literature reveals growing scholarly interest in examining quality of life among patients with diabetes from various perspectives, including correlational approaches linking self-management and health behaviors to quality of life, clinical approaches emphasizing multidisciplinary care, and demographic analyses exploring associations with socioeconomic and medical variables. Moreover, several studies have relied on the World Health Organization's instruments, reflecting the prominence of the health-related quality of life framework in this field.

However, a closer analytical examination indicates that most studies have predominantly adopted psychological or clinical orientations, often focusing on isolated variables or relatively limited samples. Comprehensive sociological analyses that integrate structural, environmental, and subjective dimensions remain comparatively scarce. Furthermore, within the Saudi context, empirical investigations addressing quality of life among patients with type 2 diabetes through a multidimensional and culturally grounded framework remain limited.

Given that quality of life is inherently a relative and context-dependent construct shaped by cultural values, social structures, and environmental conditions, there is a pressing need for research that adopts an integrative perspective grounded in health-related quality of life theory. The present study seeks to address this gap by providing a comprehensive sociological assessment of the physical, psychological, social, and environmental dimensions of quality of life among patients with type 2 diabetes in Makkah. In doing so, it extends beyond clinical outcome measures toward a broader understanding of sustained health and adaptive living in the context of chronic illness.

## **4. Methodology of the Study:**

### **4.1. Research Design:**

This study adopts a descriptive research design aimed at examining the status of quality of life dimensions among patients with type 2 diabetes. The social survey method using a sample was employed as it is appropriate for quantitatively investigating social phenomena within large populations.

### **4.2. Population and Sample:**

The study population consisted of all patients diagnosed with type 2 diabetes and registered at the diabetes and endocrinology centers affiliated with Hira General Hospital and Al Noor Specialist Hospital in Makkah. According to official 2024 statistics, 22,168 patients were registered at Hira General Hospital and 24,844 at Al Noor Specialist Hospital, totaling 47,012 patients.

A stratified random sample of 381 patients was selected, including 180 participants from Hira General Hospital and 201 from Al Noor Specialist Hospital.

### 4.3. Data Collection Instrument

Data were collected using a structured questionnaire consisting of two sections:

- Demographic information
- A 32-item scale measuring four dimensions (8 items per dimension): physical, psychological, social, and environmental.

Responses were measured using a three-point Likert scale (Always Applies – Applies to Some Extent – Does Not Apply).

### 4.4. Validity and Reliability

To ensure the validity and reliability of the research instrument (questionnaire), several statistical procedures were conducted as follows:

#### 4.4.1 Face Validity

Face validity was established by presenting the initial version of the questionnaire to a panel of experts specialized in sociology and health-related fields. The experts were asked to evaluate the clarity of the items, their linguistic formulation, their relevance to the domains they belong to, and their suitability for achieving the study objectives. Based on their feedback, necessary modifications were made, including rewording, deletion, and addition of certain items. Accordingly, the final version of the instrument was considered to have achieved acceptable face validity.

#### 4.4.2 Internal Consistency Validity

After establishing face validity, the questionnaire was administered to a pilot sample of 30 patients (after excluding five incomplete responses). Pearson correlation coefficients were calculated between each item and the total score of the domain to which it belongs. The results are presented in Table 1.

**Table 1. Pearson Correlation Coefficients Between Each Item and Its Corresponding Domain Score**

Item	Physical Domain	Item	Psychological Domain	Item	Social Domain	Item	Environmental Domain
1	0.771*	1	0.463*	1	0.771*	1	0.463*
2	0.795*	2	0.784*	2	0.795*	2	0.784*
3	0.762*	3	0.516*	3	0.762*	3	0.516*

4	0.613*	4	0.417*	4	0.613*	4	0.417*
5	0.736*	5	0.371*	5	0.736*	5	0.371*
6	0.588*	6	0.616*	6	0.588*	6	0.616*
7	0.780*	7	0.706*	7	0.780*	7	0.706*
8	0.594*	8	0.632*	8	0.594*	8	0.632*

\* Statistically significant at  $p < 0.05$

All correlation coefficients were statistically significant at the 0.05 level, indicating a high degree of internal consistency validity. This confirms that all items and domains of the questionnaire adequately measure the intended constructs.

#### 4.4.3. Construct Validity

Construct validity was examined by calculating Pearson correlation coefficients between each domain score and the total questionnaire score. The results are presented in Table 2.

**Table 2. Pearson Correlation Coefficients Between Each Domain and the Total Questionnaire Score**

Domain	Correlation with Total Score
Physical Domain	0.835*
Psychological Domain	0.630*
Social Domain	0.723*
Environmental Domain	0.596*

\* Statistically significant at  $p < 0.05$

As shown in Table 2, correlation coefficients ranged from 0.596 to 0.835 and were statistically significant at the 0.05 level, indicating satisfactory construct validity of the instrument.

#### 4.4.4 Reliability of the Instrument

To assess the reliability of the questionnaire, Cronbach’s alpha coefficients were calculated based on responses from the pilot sample. The results are shown in Table 3.

**Table 3. Cronbach’s Alpha Coefficients for Each Domain and the Overall Scale**

Domain	Number of Items	Cronbach’s Alpha
Physical Domain	8	0.822
Psychological Domain	8	0.821

Social Domain	8	0.809
<b>Environmental Domain</b>	8	0.895
<b>Overall Scale</b>	32	0.880

As shown in Table 3, Cronbach's alpha values ranged between 0.809 and 0.895 across the four domains, while the overall reliability coefficient reached 0.880. These results indicate a high level of internal consistency and reliability of the instrument.

## 5. Results and Discussion:

### 5.1. Sample Characteristics

To describe the demographic and background characteristics of the study sample, frequencies and percentages were calculated.

#### 5.1.1 Distribution by Hospital Affiliation

The distribution of participants according to hospital affiliation is presented in Table 4.

**Table 4. Distribution of Participants by Hospital Affiliation**

Hospital	Frequency	Percentage
Al Noor Specialist Hospital	201	52.8%
Hira General Hospital	180	47.2%
<b>Total</b>	381	100%

As shown in Table 4, 52.8% of the participants were recruited from Al Noor Specialist Hospital, while 47.2% were from Hira General Hospital. This distribution reflects the proportional size of the registered diabetic population in both centers.

#### 5.1.2 Distribution by Gender

The distribution of participants according to gender is shown in Table 5.

**Table 5. Distribution of Participants by Gender**

Gender	Frequency	Percentage
Male	212	55.6%
Female	169	44.4%
<b>Total</b>	381	100%

Table 5 indicates that males constituted 55.6% of the sample, whereas females represented 44.4%.

### 5.1.3. Distribution by Age Group:

Table 6 presents the distribution of participants according to age group.

**Table 6. Distribution of Participants by Age Group**

Age Group	Frequency	Percentage
Less than 30 years	9	2.4%
30–39 years	20	5.2%
40–49 years	75	19.7%
50–59 years	119	31.2%
60 years and above	158	41.5%
<b>Total</b>	<b>381</b>	<b>100%</b>

As shown in Table 6, the majority of participants (41.5%) were aged 60 years and above, followed by those aged 50–59 years (31.2%).

### 5.1.4 Distribution by Monthly Expenditure on Diabetes Care

The distribution of participants according to monthly expenditure on diabetes care is presented in Table 7.

**Table 7. Distribution of Participants by Monthly Expenditure on Diabetes Care**

Monthly Expenditure (SAR)	Frequency	Percentage
Less than 1000	128	33.6%
1000–1999	198	52.0%
2000–2999	44	11.5%
3000 and above	11	2.9%
<b>Total</b>	<b>381</b>	<b>100%</b>

Table 7 shows that more than half of the participants (52.0%) reported spending between 1000 and 1999 SAR monthly on diabetes care.

### 5.1.5. Distribution by Duration of Complications After Diagnosis

Table 8 presents the distribution of participants according to the duration between diagnosis and the onset of complications.

**Table 8. Distribution of Participants by Duration of Complications**

Duration (Years)	Frequency	Percentage
Less than 10	79	20.7%

10–19	222	58.3%
20–29	54	14.2%
30 and above	26	6.8%
<b>Total</b>	<b>381</b>	<b>100%</b>

As shown in Table 8, the majority of participants (58.3%) reported that complications developed between 10 and 19 years after diagnosis.

### 5.1.6 Distribution by Motivating Symptoms for Seeking Treatment

The motivating factors for seeking treatment are presented in Table 9.

**Table 9. Motivating Symptoms for Seeking Treatment**

Motivating Symptom	Frequency	Percentage
Inability to perform daily activities easily	293	26.6%
Medical advice to regulate blood glucose	183	16.6%
Diagnosis discovery	148	13.4%
Recurrent disease episodes	134	12.2%
Physical appearance changes	122	11.1%
Supportive environment	2	0.2%
Other	219	19.9%
<b>Total Responses</b>	<b>1101</b>	<b>100%</b>

As shown in Table 9, the inability to perform daily activities was the most frequently reported motivating factor for seeking treatment (26.6%).

## 5.2. Results Related to the First Research Question

The first research question sought to examine the status of quality of life dimensions among patients with type 2 diabetes in diabetes and endocrinology centers in Makkah. To answer this question, means and standard deviations were calculated for the four dimensions of quality of life (physical, psychological, social, and environmental). The results are presented in Table 10.

**Table 10. Means and Standard Deviations of Quality of Life Dimensions Among Patients with Type 2 Diabetes**

No.	Dimension	Mean	Percentage	Std. Deviation	Response Level	Rank
1	Physical	1.70	35%	0.459	Applies to Some Extent	4
2	Psychological	2.20	60%	0.257	Applies to Some Extent	2
3	Social	2.06	53%	0.229	Applies to Some Extent	3

4	Environmental	2.45	72%	0.444	Always Applies	1
—	Overall Quality of Life	2.10	55%	0.251	Applies to Some Extent	—

As shown in Table 10, the overall mean score for quality of life was 2.10, which falls within the second category of the three-point Likert scale (1.67–2.33), indicating that quality of life among patients with type 2 diabetes generally “applies to some extent.”

Among the four dimensions, the environmental dimension ranked first ( $M = 2.45$ ), with a response level of “always applies.” The psychological dimension ranked second ( $M = 2.20$ ), followed by the social dimension ( $M = 2.06$ ), while the physical dimension ranked last ( $M = 1.70$ ).

### - Discussion of the First Research Question

The prominence of the environmental dimension may reflect the structured healthcare environment and the accessibility of diabetes-related services within Saudi Arabia. Previous reports have highlighted the expansion of specialized diabetes and endocrinology centers across the Kingdom, as well as national initiatives aimed at improving quality of life and public health infrastructure (National Unified Platform, 2022).

However, the moderate levels observed in the psychological, social, and physical dimensions may be attributed to the chronic nature of diabetes, which requires lifelong management and may impose emotional, physical, and social burdens on patients. This finding aligns with Prajapati et al. (2017), who reported that patients with diabetes generally experience lower quality of life compared to healthy individuals. Furthermore, the overall moderate level of quality of life supports the health-related quality of life framework, which emphasizes the interaction between subjective perceptions and objective health conditions. As noted by Khalaf et al. (2022), evaluating chronic illness outcomes should extend beyond clinical indicators to include psychological and social well-being.

### 5.3. Physical Dimension of Quality of Life

To address the second research question, eight items were used to assess the physical dimension of quality of life among patients with type 2 diabetes. Means and standard deviations were calculated, and the results are presented in Table 11.

**Table 11. Means and Standard Deviations of the Physical Dimension Items**

No.	Item	Mean	Percentage	Std. Deviation	Response Level	Rank
5	I take my prescribed medication regularly	2.28	64%	0.634	Applies to Some Extent	1

6	I find sufficient time to care for my health condition	2.20	60%	0.622	Applies to Some Extent	2
8	I sleep adequately without daily interruption due to diabetes episodes	2.05	52%	0.526	Applies to Some Extent	3
7	I take proper foot care (drying, moisturizing, cotton socks)	1.56	28%	0.764	Does Not Apply	4
2	I record blood glucose levels before meals	1.46	23%	0.638	Does Not Apply	5
1	I eat main meals at scheduled times	1.43	22%	0.676	Does Not Apply	6
3	I record blood glucose levels two hours after meals	1.34	17%	0.543	Does Not Apply	7
4	I monitor blood glucose during exercise to avoid episodes	1.30	15%	0.610	Does Not Apply	8
—	Overall Physical Dimension	1.70	35%	0.459	Applies to Some Extent	—

As shown in Table 11, the overall mean score for the physical dimension was 1.70, which falls within the second category of the three-point Likert scale (1.67–2.33), indicating that the physical dimension “applies to some extent.”

The highest-ranked item was regular medication adherence ( $M = 2.28$ ), followed by allocating sufficient time for health care ( $M = 2.20$ ). In contrast, items related to self-monitoring behaviors—such as recording blood glucose levels and monitoring during exercise—received the lowest mean scores, indicating limited engagement in comprehensive self-care practices.

The results reveal variation in participants’ responses, with mean scores ranging from 1.30 to 2.28, spanning both the “does not apply” and “applies to some extent” categories.

### ***- Discussion of the Physical Dimension***

The moderate overall score of the physical dimension may reflect partial adherence to medical treatment accompanied by limited behavioral self-management practices. While medication adherence appears relatively higher, lifestyle-related practices such as regular glucose monitoring, dietary regulation, and preventive foot care show lower levels of commitment.

This pattern may suggest that diabetes management among some patients remains predominantly medication-oriented rather than behavior-oriented. Effective management of chronic illnesses such as diabetes requires not only pharmacological treatment but also sustained lifestyle modifications and active self-care behaviors.

This finding is consistent with Gounas (2022), who reported that higher levels of health behavior are associated with improved quality of life among diabetic patients.

#### 5.4. Psychological Dimension of Quality of Life

To address the third research question, eight items were used to assess the psychological dimension of quality of life among patients with type 2 diabetes. Means and standard deviations were calculated, and the results are presented in Table 12.

**Table 12. Means and Standard Deviations of the Psychological Dimension Items**

No	Item	Mean	Percentage	Std. Deviation	Response Level	Rank
4	I remind myself of the spiritual reward of patience to relieve pain	2.94	97%	0.280	Always Applies	1
3	I feel loved and cared for within my family	2.76	88%	0.524	Always Applies	2
7	I feel bored due to frequent medication intake	2.67	83%	0.519	Always Applies	3
6	I feel exhausted from long-term disease management	2.65	83%	0.535	Always Applies	4
2	I am satisfied with my physical appearance	2.15	57%	0.865	Applies to Some Extent	5
8	I accept others' criticism regarding my lifestyle as a diabetic patient	1.70	35%	0.769	Applies to Some Extent	6
1	I feel comfortable despite being restricted by a special health regimen	1.50	25%	0.698	Does Not Apply	7
5	I am motivated to learn how to calculate carbohydrate intake	1.27	13%	0.590	Does Not Apply	8
—	Overall Psychological Dimension	2.20	60%	0.257	Applies to Some Extent	—

As shown in Table 12, the overall mean score for the psychological dimension was 2.20, which falls within the second category of the three-point Likert scale (1.67–2.33), indicating that the psychological dimension “applies to some extent.”

The highest mean score was observed for the item related to spiritual coping ( $M = 2.94$ ), followed by feeling loved and supported within the family ( $M = 2.76$ ). Conversely, lower mean scores were recorded for items related to lifestyle acceptance and motivation to learn carbohydrate management.

The responses showed considerable variation, with mean scores ranging from 1.27 to 2.94, spanning all three Likert categories.

### **- Discussion of the Psychological Dimension**

The findings suggest that while participants demonstrate strong spiritual coping mechanisms and family support, challenges remain in areas related to lifestyle adaptation and proactive disease management. Spiritual resilience appears to play a significant role in psychological adjustment, which aligns with previous literature emphasizing the role of religious coping in managing chronic illness stress.

At the same time, the presence of boredom and exhaustion related to long-term disease management reflects the psychological burden associated with chronic conditions. The American Diabetes Association (2023) highlights that patients with diabetes often experience persistent emotional distress due to the ongoing demands of disease management. These results also support the psychological perspective of quality of life, which emphasizes subjective perceptions and emotional responses as central components of well-being. As indicated by Grio (2022), psychological support contributes significantly to improving quality of life among patients with diabetes.

### **5.5. Social Dimension of Quality of Life**

To address the fourth research question, eight items were used to assess the social dimension of quality of life among patients with type 2 diabetes. Means and standard deviations were calculated, and the results are presented in Table 13.

**Table 13. Means and Standard Deviations of the Social Dimension Items**

No.	Item	Mean	Percentage	Std. Deviation	Response Level	Rank
5	I offer help to others	2.80	90%	0.424	Always Applies	1
7	I lose control of blood glucose levels during social occasions	2.67	83%	0.577	Always Applies	2

8	Conflicts with others increase due to my illness	2.49	74%	0.671	Always Applies	3
6	I receive support during family crises	2.43	71%	0.639	Always Applies	4
3	I perform my daily activities easily	1.94	47%	0.575	Applies to Some Extent	5
2	I recognize the benefits of exercise for diabetes	1.70	35%	0.784	Applies to Some Extent	6
1	I practice appropriate physical exercise	1.31	16%	0.619	Does Not Apply	7
4	I attend awareness activities organized by the center	1.18	9%	0.453	Does Not Apply	8
—	Overall Social Dimension	2.06	53%	0.229	Applies to Some Extent	—

As shown in Table 13, the overall mean score of the social dimension was 2.06, which falls within the second category of the three-point Likert scale (1.67–2.33), indicating that the social dimension “applies to some extent.”

The highest mean score was observed for helping others ( $M = 2.80$ ), followed by difficulty maintaining glycemic control during social occasions ( $M = 2.67$ ). In contrast, participation in awareness activities ( $M = 1.18$ ) and practicing appropriate physical exercise ( $M = 1.31$ ) received the lowest mean scores. The results reveal substantial variation across items, with mean scores ranging from 1.18 to 2.80, covering all three Likert response levels.

### ***- Discussion of the Social Dimension***

The moderate overall level of the social dimension suggests that patients remain socially active and engaged, yet face challenges in balancing disease management with social participation. While strong indicators of social contribution and perceived support were observed, difficulties in glycemic control during social gatherings and increased interpersonal tensions reflect the complex interaction between chronic illness and social life. From a sociological perspective, quality of life within the social framework emphasizes belonging, social integration, and access to supportive networks. Social interaction fulfills individuals’ needs for acceptance and recognition within their community. The presence of both supportive relationships and illness-related conflicts highlights the dual role of social environments in either buffering or amplifying the burden of chronic illness.

Empirical evidence also supports the importance of social capital in improving quality of life among patients with type 2 diabetes, as previous research has demonstrated a positive and statistically significant relationship between social capital and quality of life.

### 5.6. Environmental Dimension of Quality of Life

To address the fifth research question, eight items were used to assess the environmental dimension of quality of life among patients with type 2 diabetes. Means and standard deviations were calculated, and the results are presented in Table 14.

**Table 14. Means and Standard Deviations of the Environmental Dimension Items**

No.	Item	Mean	Percentage	Std. Deviation	Response Level	Rank
5	Authorized healthcare centers are located near my residence	2.82	91%	0.477	Always Applies	1
3	Medication delivery services are available in my neighborhood	2.80	90%	0.494	Always Applies	2
6	Healthcare services are easily accessible near my residence	2.61	81%	0.617	Always Applies	3
1	I live in adequate and healthy housing	2.58	79%	0.596	Always Applies	4
4	I live in a clean neighborhood	2.42	71%	0.766	Always Applies	5
8	I have transportation that allows me to attend medical appointments	2.35	67%	0.828	Always Applies	6
2	There is a suitable place for exercise in my neighborhood	2.15	58%	0.915	Applies to Some Extent	7
7	Recreational facilities suitable for my health are available nearby	1.85	43%	0.619	Applies to Some Extent	8
—	Overall Environmental Dimension	2.45	72%	0.444	Always Applies	—

As shown in Table 14, the overall mean score for the environmental dimension was 2.45, which falls within the third category of the three-point Likert scale (2.34–3.00), indicating that the environmental dimension “always applies.” The highest mean scores were related to the availability and accessibility of healthcare services and medication delivery, whereas lower scores were observed

for recreational facilities and exercise-friendly environments. Mean values ranged from 1.85 to 2.82, indicating generally favorable environmental conditions with some variability across specific aspects.

### ***- Discussion of the Environmental Dimension***

The high overall score of the environmental dimension suggests that participants benefit from supportive structural and infrastructural conditions that facilitate access to healthcare services. The strong presence of nearby authorized healthcare centers, medication delivery services, and transportation accessibility reflects the role of healthcare infrastructure in shaping patients' quality of life. From a sociological perspective, the environmental dimension represents the structural context within which individuals manage chronic illness. Social theory emphasizes that quality of life is not solely determined by individual factors but also by the availability of community-level resources, public services, and living conditions that enable individuals to meet their needs effectively. Previous studies have also demonstrated that social and community support structures contribute significantly to improving quality of life among patients with chronic illnesses, including type 2 diabetes.

### **6. Conclusion:**

This study examined the multidimensional nature of quality of life among patients with type 2 diabetes attending diabetes and endocrinology centers in Makkah. The findings indicate that overall quality of life was perceived at a moderate level, suggesting that patients experience a partial but not comprehensive sense of well-being in managing their condition.

The environmental dimension emerged as the most prominent domain, reflecting the structural strength of healthcare accessibility and service availability. This finding underscores the importance of institutional and infrastructural support in shaping lived experiences of chronic illness.

In contrast, the physical dimension recorded the lowest level among the four domains, revealing gaps in sustained self-care behaviors despite relatively acceptable levels of medication adherence. This suggests that disease management remains partially biomedical rather than fully integrative.

The psychological and social dimensions presented a mixed pattern. While participants demonstrated strong spiritual coping and perceived family support, they also reported fatigue, boredom, and social strain associated with prolonged disease management. These findings highlight the complex interaction between personal resilience and the cumulative burden of chronic illness.

Taken together, the results affirm that quality of life among patients with diabetes cannot be reduced to glycemic control alone. Rather, it is shaped by the interplay between structural conditions,

behavioral practices, emotional adaptation, and social integration. This reinforces the relevance of the health-related quality of life framework as a comprehensive lens for understanding chronic disease experiences.

## 7. Recommendations

In light of the findings, the study proposes the following:

1. Developing structured educational interventions that move beyond medication adherence to emphasize sustainable self-management behaviors.
2. Integrating psychological counseling services into routine diabetes care to address long-term emotional fatigue and adaptive stress.
3. Encouraging active participation in community-based awareness and peer-support initiatives to strengthen social engagement.
4. Promoting family-inclusive care models that reinforce supportive environments for chronic disease management.
5. Adopting an integrative care strategy that aligns biomedical treatment with psychosocial and environmental determinants of health.
6. Expanding sociological research within the Saudi context to further explore how cultural values, social capital, and environmental infrastructure interact to shape quality of life among individuals living with chronic illness.

## 8. Acknowledgment

The authors would like to express their sincere appreciation to all participants who contributed to this study. The authors also declare that there is no conflict of interest related to this research. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

## 9. References

- Al Amin, B. M. (2020). *Quality of life among patients with diabetes* (Master's thesis). Abdelhamid Ben Badis University, Mostaganem, Algeria.
- Al Anzi, S. G. (2023). Quality of life: Types, dimensions, indicators, and explanatory approaches. *Journal of Social Sciences Future*, 1(15), 65–78.
- Al Dawish, M., Robert, A., Braham, R., Al Hayek, A., Al Saeed, A., Ahmed, R., & Al Sabaan, F. (2016). Diabetes mellitus in Saudi Arabia: A review of the recent literature. *Current Diabetes Reviews*, 12(4), 359–368.

- Al Ghazali, M. N. (2012). *Nutrition for diabetic patients*. Anglo Egyptian Library.
- Alharbi, M., et al. (2024). Burden of illness of type 2 diabetes mellitus in the Kingdom of Saudi Arabia: A five-year longitudinal study. *Advances in Therapy*, 41, 1120–1150.
- Al Qawasmi, J. (2020). Measuring quality of life in urban areas: Toward an integrated approach. *International Journal of Environmental Sciences & Natural Resources*, 25(2), 67–74.
- American Diabetes Association. (2023). *About diabetes*. <https://diabetes.org>
- Bayoumi, M. A. (1996). *Religious sociology and problems of the Islamic world* (3rd ed.). Dar Al-Maaref Al-Jamiya.
- Bouti, S., & Barika, Y. (2020). A theoretical perspective on quality of life: Concept and trends.
- Cardona, L. A. (2010). *Understanding quality of life in older adults* (Master's thesis). University of North Texas.
- Clark, K. K. (2024). Implementation of diabetic retinopathy screening in adult patients with type 2 diabetes in a primary care setting. *Clinical Diabetes*, 42(2), 223–231.
- Conner, B. L. (2016). *A systematic review of quality of life assessment tools for adults with intellectual and developmental disabilities* (Master's thesis). The University of New Mexico.
- Grew, S. (2022). The role of medical, psychological, and social support in improving quality of life among diabetic patients: A case study. *Insaniyah Journal for Research and Studies*, 2(13).
- International Diabetes Federation. (2023). *What is diabetes?* <https://idf.org/about-diabetes/what-is-diabetes/>
- John, R., et al. (2019). Evaluation of quality of life in type 2 diabetes mellitus patients using quality of life instrument for Indian diabetic patients: A cross-sectional study. *Journal of Mid-life Health*, 10(2), 81–88.
- Kerce, E. W. (1992). *Quality of life: Meaning, measurement, and models*. Navy Personnel Research and Development Center.
- Keyvanara, M., et al. (2018). The relationship between social capital and quality of life among patients referring to diabetes centers in Isfahan, Iran. *Journal of Diabetes Research*, 1–6.
- Maamria, B. (2020). Quality of life: Definition, determinants, manifestations, and dimensions. In *Proceedings of the National Forum on Quality of Life and Sustainable Development in Algeria* (pp. 15–26).
- Ministry of Health, Kingdom of Saudi Arabia. (2024). *Health awareness days 2024*. <https://www.moh.gov.sa>

- Ministry of Health, Kingdom of Saudi Arabia. (1441–1445H). *Strategic plan 1441–1445H*.
- Ministry of Health, Syrian Arab Republic. (2016). *Educational guide for diabetes in primary care and acute complications*.
- National Institute of Diabetes and Digestive and Kidney Diseases. (2017). *Type 2 diabetes*. <https://www.niddk.nih.gov>
- Nguyen, T. K., Hoa, N. P., Duc, D. M., & Wens, J. (2021). Health-related quality of life and associated factors among patients with type II diabetes mellitus. *Health Psychology Open*, 1–9.
- Prajapati, V. B., et al. (2017). Assessment of quality of life in type II diabetic patients using the modified diabetes quality of life questionnaire. *Brazilian Journal of Pharmaceutical Sciences*, 53(4), 1–9.
- Qounas, I. (2022). Health behavior and quality of life among patients with diabetes: A clinical study.
- Rocha, R. B., et al. (2020). Self-care in adults with type 2 diabetes mellitus: A systematic review. *Current Diabetes Reviews*, 16(6), 598–607.
- Rossi, M., et al. (2010). Diabetes interactive diary: A new telemedicine system enabling flexible diet and insulin therapy while improving quality of life. *Diabetes Care*, 33(1), 109–115.
- Saudi Vision 2030. (2018). *Quality of Life Program: Delivery plan*.
- Saudi Vision 2030. (2020). *Health Sector Transformation Program: Delivery plan*.
- Surmach, M. J., & Piecewicz-Szczęśna, H. (2014). Evaluation of quality of life of children in terms of health as an interdisciplinary problem. *Polish Journal of Public Health*, 124(3), 148–151.
- Unified National Platform. (2022). *Healthcare services*. <https://www.my.gov.sa>
- World Health Organization. (2012). *WHOQOL user manual*. World Health Organization.
- Zitouni, A., & Rebhouh, N. (2024). *Self-management and its relationship to quality of life among patients with type 2 diabetes* (Master's thesis). Ziane Achour University, Djelfa, Algeria.

Copyright © 2026 by Khairiya Sami Ngoum, Prof. Ayedh Saad Al-Shahrani, and AJRSP. This is an Open-Access Article Distributed under the Terms of (CC BY NC)

Doi: <https://doi.org/10.52132/Ajrsp.e.2026.83.2>